

# IS NERVE PAIN DISRUPTING YOUR LIFESTYLE?

There are Answers & Solutions



A Guide to  
Understanding  
Lower Extremity  
Nerve Pain

HARVEY R. DANCIGER, DPM

# Is Nerve Pain Disrupting Your Lifestyle?

There are Answers & Solutions

Dr. Harvey Danciger

**Harvey R. Danciger, DPM**

74-000 Country Club Drive, Suite A-2

Palm Desert, CA 92260

(760) 568-0108 Office • (760) 568-5110 Fax

**[www.CoachellaValleyPodiatrist.com](http://www.CoachellaValleyPodiatrist.com)**

Copyright © 2015 by Dr. Harvey Danciger. All rights reserved. No part of this book may be used or reproduced in any manner whatsoever without written permission of the author. No expressed or implied guarantees have been made or are made by the author or publisher. Individual results may vary. Neither author nor publisher accepts any liability or responsibility to any person with respect to any loss or damage alleged to have been caused by the information in this book. Always seek professional medical advice.

# Table of Contents

|   |    |
|---|----|
| Why I Wrote This Book .....                                     | 4  |
| What Is Lower Extremity Nerve Pain? .....                       | 6  |
| Common Nerve Conditions of the Foot and Ankle.....              | 7  |
| What Are the Symptoms of Lower Extremity Nerve Pain? .....      | 11 |
| What Causes Lower Extremity Nerve Pain?.....                    | 12 |
| What Can a Podiatrist Do About Lower Extremity Nerve Pain?..... | 13 |
| How Does a Podiatrist Treat Other Conditions?.....              | 17 |
| What Can You Do to Keep Your Feet Healthy? .....                | 24 |
| When to Call a Podiatrist .....                                 | 25 |
| What to Do If You Are In Severe Pain .....                      | 26 |
| Final Thoughts .....  | 27 |

## Why I Wrote This Book

As a podiatrist, I specialize in the care and treatment of the feet and ankles. In the last decade, I've seen firsthand that the incidence of lower extremity nerve pain has increased as our general population continues to age.

**As many as 28-32 million Americans suffer from lower extremity nerve pain.**

Lower extremity nerve pain is experienced by people of all ages and demographic backgrounds. It most often occurs in middle age or later, although it might be the result of an injury that happened earlier in life. It is sometimes related to an underlying metabolic disease such as diabetes.

There are a number of causes of lower leg and foot pain, and I will cover them in greater detail later in this book. What concerns me most is that often, this common problem goes undiagnosed because symptoms can be so varied and because the onset of symptoms is very slow.

Frequently, the people that come in to see me do so after the symptoms have become so severe that treatment options are limited. The paradox of this condition is that it often goes undiagnosed, yet early treatment is the key to successfully eliminating the problem.

After noticing all of these issues, I decided to create a publication for you. I wrote it so that you could understand the symptoms and causes of lower extremity nerve pain, and the treatments available to you. I'm very excited about the new treatment options for nerve pain that are available to podiatrists, including the use of minimally invasive surgery, and I want you to know that we have the very latest technologies available in our office. Dr. Harvey Danciger is trained to perform nerve decompression surgery.

## Is Nerve Pain Disrupting Your Lifestyle?

If after reading this book you think you might be helped by seeing a podiatrist, I hope you consider Harvey R. Danciger, DPM. My staff and I will do everything we can to treat your condition, eliminate your pain, and make you feel at home.

Dedicated to Your Health,

*Dr. Harvey Danciger*

## **What is Lower Extremity Nerve Pain?**

Many of the cases of lower extremity nerve pain are classified as peripheral neuropathy. This is a disorder of the nerves that connect the spinal cord to the muscles, skin, and organs. The peripheral nerves control your sense of touch, the way you feel pain and temperature, and your muscle strength. Most of the time, it starts with a loss of feeling in the toes, and then moves into the limbs causing pain and loss of feeling in the feet and legs. Patients with peripheral neuropathy may have the same symptoms in their fingers and hands as well.


Another type of lower extremity nerve pain is caused by nerve entrapment. This is commonly known as a pinched nerve and causes burning pain, shooting or sensitivity to the top or bottom of the foot. Nerve entrapments may be caused by trauma to the foot or repetitive compression like wearing too tight shoes or running in shoes that do not fit properly.

Nerve injury is the third type of lower extremity nerve pain. Nerve injury is also caused by prolonged repetitive compression of the nerves or trauma to the foot or lower leg.

# Common Nerve Conditions of the Foot and Ankle


## Tarsal Tunnel Syndrome

At the inside of your ankle, a nerve divides into branches, and crosses from the leg into the foot through a series of tunnels. This nerve is called the Tibial Nerve, and it branches into the medial and lateral plantar nerves, and the calcaneal nerves.



When the ankle moves, the nerve gets pressed. This can happen during normal walking when there is too much motion in the foot causing the ankle to roll inward. When the nerve gets pressed, blood flow to the nerve slows, blood flow decreases, and the nerve sends a message of numbness and tingling, or buzzing of the foot/toes. Most

patients have similar complaints, and you should seek treatment when your feet or toes are numb most of the day, your foot or feet disturb your sleep every night, you have trouble feeling the floor or the car's gas or brake pedals, you begin to lose your balance, or there is pain associated with the numbness.






### Compression of the Deep Fibular (Peroneal) Nerve

Another anatomic site of entrapment is at the level of the dorsum of the foot, along the distribution of the deep fibular (peroneal) nerve sensory branch.

There is a known site of compression of the deep fibular nerve by the extensor hallucis brevis tendon at the metatarsal cuneiform joint, the joint at the peak of the arch of the foot. This is often a complaint in men and women who wear shoes that increase the pressure over this nerve. Common complaints are tingling, numbness, and burning

into the first and second toes, relieved by removal of the shoe and rubbing the dorsal arch of the foot. The deep fibular nerve can also be compressed at the front of the ankle and can cause pain on the top of the foot. Both these areas can be compressed at the same time.



### Compression of the Common Fibular (Peroneal) Nerve

Another anatomic site of compression is at the level of the outside of the leg near the knee. This is the nerve that can become compressed if you have been sitting with your legs crossed and you get up and your foot is asleep. The nerve is close under the skin as it courses around the bone on the outside


part of the leg below the knee. This nerve is the nerve mostly responsible for a condition called drop foot.

### Compression of the Sural Nerve, Superficial Fibular (Peroneal) Nerve, Saphenous Nerve

These nerves can also be compressed in the leg. The Sural Nerve goes down the outside or lateral aspect of the leg into the foot and the Superficial Fibular Nerve becomes superficial at the skin in the lower 1/3rd of the outside of the leg about 10 centimeters up from the bone on the outside of the ankle. The Saphenous Nerve is located on the medial or inside part of the leg. This can cause any of the symptoms of neuropathy or nerve pain in these areas or problems in the foot.

### Compression of the Proximal Tibial Nerve at the Soleal Sling

This nerve can become compressed in the back of the leg just below the back of the knee. Symptoms can vary among patients. Oftentimes, the patient is unable to rise up on the ball of the foot and toes using one leg about 10 times, or there can be a feeling of weakness or tightness of the calf muscles when comparing one leg to the other. Pressing into the muscle behind and just below the knee may reproduce tenderness or radiating pains which can go into the leg or foot.



### Interdigital Neuroma

Compression of the interdigital sensory nerve to the inner space of the toes in the foot is a very common complaint. Most people refer to the burning, numbness, cramping, and ball of foot pain as a Morton's Neuroma. Although this may be correct, a Morton's Neuroma only describes a compression of the interdigital nerve of the third inner space, affecting the third and fourth toes only. These symptoms are most common while wearing a closed toe shoe, a high heeled shoe, and during endurance exercise, like running or cycling. Nerve compression in the toe can occur in all the inner spaces, but the third inner space is the most common. A thorough history and physical examination will determine the presence of this condition, and only after a definitive diagnosis is made will a treatment regimen be discussed. Treatments include patient education, shoe modification, custom orthotics, laser treatments, oral anti-inflammatory medications, water soluble steroid injections, and surgical decompression of the inner space. Some antidotal evidence on the use of alcohol sclerosis agents may be discussed. Ultimately, if conservative and palliative therapies fail to mitigate the pain, surgical decompression or removal of the sensory nerve may be necessary.



## **What Are the Symptoms of Lower Extremity Nerve Pain?**

Symptoms of lower extremity nerve pain can develop slowly over time. They also vary widely from patient to patient, causing diagnosis to often go undetected. Some people with nerve damage show no symptoms at all.

Symptoms of peripheral neuropathy may include pain, burning, tingling, pins and needles, electric shocks, numbness, cramping, and tightness. Some people say they feel like they are wearing socks when they are barefoot. Others describe a sensation of ants crawling on their skin. Insensitivity to heat or cold can also be a symptom. People sometimes describe a feeling of restless leg. A feeling of heaviness in your legs can be a symptom, as well as a loss of balance. Muscle weakness or paralysis may also be a symptom. Any one of these symptoms can result in difficulty in walking, causing a wider stance and a less rhythmic or fluid gait. Other symptoms are loss of feeling in the feet, cramping in the feet, curling of the toes, weakness, and sharp, shooting pains in the legs and feet. The pain is often worse at night, and some patients find relief with walking.

Symptoms of nerve entrapment include: lower back pain, continued pain after a knee replacement surgery, continued pain after an injury to the foot, ankle or leg, burning, numbness, a feeling of tightness in the leg or ankle, weakness, or restless leg syndrome. In advanced cases, a gait abnormality such as drop foot, which is due to weakness of the muscles in the anterior (front) portion of the leg, is caused by damage to the nerve located on the outside of the knee.

Other signs of a possible nerve problem can be thickened toenails, and/or thin, dry skin. This is because the nerves can also affect the autonomic nervous system.

## What Causes Lower Extremity Nerve Pain?

Peripheral neuropathy is nerve damage that generally results from a systemic disease. However, a full 30% of peripheral neuropathy cases are referred to as idiopathic, meaning they come from unknown causes.

The most common form of neuropathy is from **diabetes**; up to 30% of the cases of peripheral neuropathy are generally caused by this disease. The fact is that having your blood sugar too high over a long time can cause nerve swelling leading to nerve damage. The swelling causes the nerve to enlarge; therefore it becomes compressed as it courses down the leg through different leg compartments.

There are other diseases and conditions that can cause peripheral neuropathy. Liver or Kidney problems can lead to blood toxicity which can damage nerves. Infectious and inflammatory diseases such as Guillain-Barré Syndrome, Shingles, Lyme disease, Liver or Kidney disease, or HIV can damage nerves as well. In some cases, chemotherapy, exposure to toxic substances, heavy metal toxicity, recreational drug use, leprosy, and alcoholism can all result in damage to the peripheral nerves.

Other causes include thyroid disorders, arthritis (rheumatoid and lupus), alcoholism, vitamin deficiencies especially Vitamin B1, B6, or B12 and, heredity. Tumors on the nerve or certain medications such as those used in chemotherapy can also be a cause. Many people today are overweight, have high cholesterol, or have high blood pressures. These people have metabolic syndrome or Syndrome X, often referred to as pre-diabetic. Their neuropathy is often an early phase of diabetic peripheral neuropathy.

As stated earlier, nerve entrapment or nerve injury can be caused by trauma, compression, knee replacement surgery, arthroscopic knee surgery, and in some cases, severe ankle sprains or foot injuries. When a nerve gets pressed, blood flow to the nerve slows, blood flow decreases, and the nerve sends a message of numbness and tingling to the brain.

## **What Can a Podiatrist Do About Lower Extremity Nerve Pain?**

Let's start at the beginning and answer the question, **“What is a Podiatrist?”**

A podiatrist is a specialist who focuses on treatment of your feet and ankles. We are quite simply “The Experts” on your feet, heels, and ankles, and should be the very first doctor you call when you experience pain or other problems in this area. In addition to lower extremity nerve pain, I've also listed many of the other problems podiatrists treat in this book.

The DPM after our name stands for Doctor of Podiatric Medicine. Most people know that podiatrists deal with foot, heel, and ankle problems, and work with seniors, or diabetics, or people with poor circulation; what most people don't know is that podiatrists treat patients of all ages. A podiatrist is also well-qualified to help children and young adults with the problems that they can experience with their feet, heels, and ankles.

Podiatrists are required, just like any other doctor, to be licensed by the state in which they practice. When medically necessary, podiatrists can perform surgery to correct or remedy problems. Before we recommend surgery, we will explore the many conservative therapies and remedies that are available for our patients, and then recommend the very best course of treatment. As medical science advances and new therapies become available to the public, podiatrists are increasingly able to offer their patients some of the most state-of-the-art care available. I'm constantly amazed at the new therapies I can offer my patients and will discuss several of them now.

## Diagnosis

The first step in treating lower extremity nerve pain is proper diagnosis. Due to poor localization of pain and a wide variety of symptoms, diagnosing peripheral neuropathy can be difficult. If you think you have any signs of this disease, it is important to schedule a comprehensive lower extremity exam as soon as possible. **The key to treating neuropathy is early detection.** During the exam, your podiatrist will check for any signs of neuropathy. This begins with the doctor asking questions about your symptoms, your medical history, including use of alcohol, risk of infection, and exposure to toxic substances. Questions will also be asked about your family's medical history related to nerve disease and diabetes.

**Anyone with diabetes should get a complete foot exam every year.**

There are 3 basic components of the Neurological Examination: Sensory Exam, Motor Exam and the Reflex Exam. The exam will continue with the doctor checking your muscle strength and sensation – the ability to feel touch, temperature, and pain. Palpation and percussion techniques at key locations of known compressions and entrapments from the knee down into the foot will be tested first to determine the overt signs of compression and neuropathy symptoms. Further tests may include a Nerve Conduction Velocities (NCV) test. In this test, the doctor can measure how well individual nerves send electrical signals from the spinal cord to the muscles. This non-painful test is very useful in diagnosing both peripheral neuropathy and nerve entrapment. Another test that may be done is called an electromyography (EMG). This is a test to measure electrical discharges made by the muscles. It helps to determine if there is any muscle weakness present due to nerve damage. A third test, called the Pressure-Specified Sensory Device, may be performed. This painless test measures how hard the skin must be pressed in order for the patient to determine if there are one or two rounded probes moving or touching the skin. This can help the doctor determine the very subtle symptoms that are due to nerve entrapment, injury, or neuropathy. Unfortunately, oftentimes the tests come back as normal findings and the mainstay of diagnosing nerve compressions is done during the physical exam. Radiographs (X-Rays), ultrasound images, or Magnetic Resonance Images (MRI) may be ordered to determine if there are space occupying or biomechanical structures affecting the nerve. Finally, blood tests may be run to find out if there are any underlying causes of neuropathy such as diabetes, thyroid disease, kidney disease, or vitamin deficiencies.

### **Treating the Cause**

The first step in treating lower extremity leg pain is treating the cause. If peripheral neuropathy is caused by diabetes, it is important to get the blood glucose levels under control. If it is caused by alcohol, the patient should stop using alcohol. If the neuropathy is determined to be caused by vitamin deficiency, it can be treated by eating a healthy diet and taking vitamin supplements. In cases of nerve entrapment, the recommendation will be a stabilization of impaired biomechanics, a change in lifestyle, or a decrease in activities to give the nerve the chance to recover and restore itself.

### **Relieving the Pain**

The next step that the doctor will take is to try to relieve the pain that you are experiencing. One treatment modality is having Laser treatments. Laser therapy is non-invasive and not painful. This has helped numerous patients decrease their symptoms of neuropathy and pain. Another way might be treatment with over-the-counter pain medicine or oral prescription medications to help with more severe pain, numbness, tingling, and weakness. There are topical treatments available also that can bring some relief of lower extremity nerve pain. Formal physical therapy might be prescribed to help increase muscle strength, build muscle control, and mobilize the nerve through a technique called “nerve gliding.” If there are areas of nerve compressions, many times there are multiple sites of nerves being compressed found on the physical examination, then diagnostic nerve blocks can be given which can help locate and isolate different nerves, helping to form a treatment plan.

### **Surgery**

When other non-operative treatments fail to get the desired results, or if results are not long lasting, then surgery may be the best option. Nerve decompression (neurolysis) surgery allows the surgeon to open the tight area through which the nerves pass by dividing the anatomical structures that crosses the nerve. As a result, damaged nerves are able to regenerate after they are able to move more freely and glide and slide within the body to start receiving proper blood flow.

This surgery is a minimally invasive procedure and is generally done on an outpatient basis, and patients are able to walk the same day. Results can include significant pain relief, increased sensation, and improvement in balance and gait.



## Is Nerve Pain Disrupting Your Lifestyle?

It is very important that nerve decompression surgery only be performed by an experienced and formally trained nerve surgeon. When performed by a well-trained surgeon, this procedure can be very gratifying and provide immediate relief of lower extremity pain. Podiatrists who are trained to do nerve decompression surgery are also skilled at diagnosing the problem through physical examination, diagnostic studies, diagnostic imaging, and sensory nerve studies.

Sometimes, if the nerve has been injured beyond repair, the nerve has to be cut and implanted into the soft tissues or muscle in the area. This will stop the pain farther from the area where the nerve has been transected because there will be no sensation to the area the nerve normally supplies. This type of procedure is not done primarily, only if other methods fail to give relief.

## How Does a Podiatrist Treat Other Conditions?

Now that we know what a podiatrist is and some of the treatments available for lower extremity nerve pain, let's take a look at some other foot problems and the treatments we can offer to reduce or eliminate your pain. It is impossible to discuss all the potential problems that can affect the health of your feet in a book of this size, but what I want to tell you about here, are the problems we see most often at our office.

### Arthritis

Over 30 million American adults report being told by a doctor that they have some type of arthritis. It is a major cause of lost work time and serious disability for many people. Although arthritis is mainly a disease of adults, children may also have it. When a patient has arthritis, it means that the cartilage, and even the lining of their joints, has become swollen and inflamed.

There are numerous types of arthritis. The reason that your feet seem to be more susceptible to arthritis than other parts of your body is that your feet have so many joints that can be affected. The odds are just stacked against your feet. In addition, your feet and ankles bear the full weight of your entire body every single day.

While there are dozens of types of arthritis, I want to point out two of the most common.

**Osteoarthritis** – The most common type of arthritis is osteoarthritis. It is seen in many people as they age, although it may begin when they are younger as a result of injury or overuse. It is often more painful in weight bearing joints such as the knee, hip, and spine than in the wrist, elbow, and shoulder joints. All joints may be affected if they are used extensively in work or sports, or if they have been damaged from fractures or other injuries.

In osteoarthritis, the cartilage covering the ends of the bones gradually wears away. In many cases, bone growths called “spurs” can

develop in osteoarthritis joints. The joint inflammation causes pain and swelling. Continued use of the joint produces pain. Some relief may be possible through rest or modified activity.

**Rheumatoid Arthritis** – Rheumatoid arthritis is a long-lasting disease that can affect many parts of the body, including the joints. Rheumatoid arthritis can affect people of all ages, even children. However, more than 70 percent of people with this disease are over 30 years old. Many joints of the body may be involved at the same time. Arthritis cannot be cured but it can be treated. The goals of treatment are to provide pain relief, increase motion, and improve strength.

### **Bunions**

Bunions are caused by your big toe joints becoming incorrectly aligned. This causes the first joint on your big toe to slant outward and the second joint then angles toward your other toes. Your joints then begin to swell causing a bump of bone on the foot that can become very painful if left untreated.

Bunions can be hereditary, but also can be aggravated by shoes that aren't a good fit. Surgery is often recommended to correct the problem. Some bunions can be treated without surgery by wearing protective pads to cushion the painful area, orthotic devices, laser treatments, injections and of course, avoiding ill-fitting shoes in the first place.

### **Diabetes**

Diabetes can affect many parts of the body, especially the feet and heels. According to the American Diabetes Association, about 15.7 million Americans (5.9 percent of the United States population) have diabetes. It is very important that a diabetic gives the feet very special care. A small problem in a healthy person could become a severe one to a diabetic.

Diabetes can affect the feet in a number of different ways. The first is infection, which is one of the most common complications of the diabetic foot. Because diabetes causes reduced immune response, a diabetic patient's ability to fight infection is decreased. Early treatment of infection is a critical component to success. If the infection is neglected, infection of the foot can cause gangrene, ulceration, osteomyelitis (bone infection), and even amputation.

With a diabetic foot, a wound as small as a blister from wearing a shoe that is too tight can cause a lot of damage. Diabetes decreases blood flow, so injuries are slow to heal. When your wound is not healing, it's at risk for infection. As a diabetic, your infections spread quickly. If you have diabetes, you should inspect your feet every day, morning and night. Look for puncture wounds, bruises, pressure areas, redness, warmth, blisters, ulcers, scratches, and cuts. Look for any changes or anything abnormal. Get someone to help you, or use a mirror. Diabetic foot care can be very complicated, and good podiatric care is an essential component of managing diabetes.

### **Here are some basic tips for the care of diabetic feet:**

- Inspect your feet every day.
- Keep your feet clean and dry.
- Always keep your feet warm.
- Take extra care drying your feet and toes after showering. Pay special attention to the space between your toes.
- Be sure to exercise. Walking is one of the best exercises for diabetics unless you have complications. If you struggle with balance, use a cane.
- Always protect your feet and legs. Never walk barefoot. Avoid hot water bottles and heating pads.
- Do not overexpose skin to the sun.
- Never use razor blades, knives, scissors, or medicated corn/wart removers.
- Look for redness, blisters, scratches, cracks between the toes, discoloration, or any other change.
- Avoid all actions that diminish circulation such as tobacco use, sitting with legs crossed, and circular elastic garters.
- Change your shoes and socks daily.
- Wear soft leather shoes that conform to the shape of your overall foot.
- Gradually “break in” new shoes to avoid blisters.
- Call us immediately if you see any changes in your feet.

### **Fungal Nails**

Fungal infections of the nail bed, matrix, or nail plate are responsible for about 50% of all cases of thick, discolored toenails. There are other problems; for example psoriasis, trauma, thyroid disease, certain vitamin deficiencies, which can cause the nail to look like a fungal toenail but are treated differently, so diagnosis is very important. There are four different types of fungal nail infections classified by the part of the nail involved. Fungal nails can be caused

by tight footwear, minor trauma caused by exercise, communal showers, improper nail care, pedicures, and diseases that influence the immune system.

Treatment for fungal nails varies by the nature of the infection and the severity. A podiatrist can detect a fungal infection early and formulate a suitable treatment plan. This can include topical or oral medication, debridement, laser therapy, and in extreme cases, surgery. Trying to solve any nail infection without the help of a podiatrist can lead to more problems.

### **Hammertoe**

Hammertoe is a flexible or rigid contraction usually affecting the second, third, fourth, or fifth toe. In this condition, the toe is bent at the middle joint, resembling a hammer. Muscle imbalance leads to a bending or “buckling” of the toe joints. These buckled or contracted positions create any number of problems within and on top of the toe deformity. It is important to treat hammertoes early. As they advance and lose flexibility, the only option for correction may be surgery. Hammertoes can cause complications such as corns or calluses at the point where they come into contact with the shoes. As with many foot problems, one of the causes of hammertoes can be improperly fitted shoes.



Podiatrists have a variety of ways to treat hammertoes, including surgery, better shoes designed with extra room for toes, corn pads, straps, cushions, and orthosis.

### **Metatarsalgia**

Metatarsalgia is foot pain in the ball of your foot which is the area between your arch and your toes. It gets its name because the pain experienced is located in the metatarsals located in this part of the foot.

In this condition, one or more of the metatarsal joints becomes inflamed and possibly painful. People often develop a callus under the affected joint. There are many causes of Metatarsalgia including injuries, arthritis, poorly fitting shoes, and working on very hard surfaces. Sometimes, changing your shoes will fix the problem. We may also recommend orthotics or implants.

### **P.A.D.**

Peripheral Arterial Disease, known as P.A.D., is the narrowing or blockage of arteries in the legs that result in poor blood flow. Just like clogged arteries in

your heart, clogged arteries in your legs mean you are at higher risk of getting a heart attack or stroke. Plaque buildup in the legs does not always cause symptoms; so many people can have P.A.D. and not know it. Symptoms, when they are present, can include pain and cramping in the legs.

Podiatrists can perform P.A.D. screening during a foot exam using a variety of methods and tests. Treatment goals are to reduce symptoms and prevent heart attack and stroke. Treatments include making lifestyle changes such as quitting smoking, exercising, lowering blood pressure, cholesterol, and blood glucose levels, and achieving a healthy weight. Medications may be prescribed to lower blood pressure and cholesterol, to prevent the formation of blood clots that could cause a heart attack or stroke, or to reduce leg pain while walking. In some advanced cases, surgery may be recommended to improve blood circulation to the legs and the ability to walk.

### **Plantar Fasciitis (Heel Pain)**

Heel pain is the most common complaint that we see in our office. Plantar Fasciitis is an inflammation in the band of tissue (the plantar fascia) that runs from the heel to the toes. This condition is most often caused by poor foot structure such as overly flat feet or high arches. It can also be caused by an injury, wearing non-supportive footwear on hard surfaces, spending long hours on your feet, or obesity. The pain from plantar fasciitis is usually a sharp, stabbing pain on the inside of the bottom of the heel that can feel like a knife sticking into your heel. Pain from plantar fasciitis is usually most severe when you first stand on your feet in the morning. It will usually subside, but can return with prolonged standing or walking, or getting up after long periods of sitting.

We offer a variety of treatments for heel pain including night splints, stretching, ice, laser treatments, anti-inflammatory medication, and orthotics. There are always advances being made in the way that heel pain is treated so you should seek the advice of a podiatrist if you are experiencing pain for longer than 4-5 days.

### **Sprains**

An ankle sprain occurs by stretching or tearing one or more ligaments on either or both sides of the ankle. Ignoring a sprain won't help it heal any faster. Ankle injuries that are serious enough to cause disabling pain should be treated by a podiatrist. Further examination may even reveal a torn ligament or bone fracture. Common treatments for sprains include rest, eleva-

tion, compression, and ice. More serious sprains may call for immobilization, casts, crutches or other walking devices.

### **Stress Fractures**

An incomplete break in the bone caused by overuse is known as a stress fracture. Symptoms can include pain, swelling, and redness. Up to 15% of all sports injuries are stress fractures. A podiatrist needs to perform an examination and look at X-rays of the injury in order to diagnose a stress fracture. Musculoskeletal ultrasound can often detect stress fractures not yet visible on x-rays. Treatments include immobilization of the foot with the use of a cast, surgical shoe, medications, and in some cases, orthotic devices to prevent further injury.

### **Warts**

Warts are caused by a virus that generally enters the body through small nicks or abrasions in the skin. When they occur on the soles of the feet, they are known as plantar warts. Due to the amount of pressure that is put on the feet in the course of a day, plantar warts can become quite painful. Teenagers between the ages of 12–16 are most commonly infected by warts, but they can occur at any age. Warts are often contracted by walking barefoot on dirty surfaces or ground. The virus thrives in warm, moist environments like showers and swimming pools.

If you suspect that you or a family member has a plantar wart, see a podiatrist to get a correct diagnosis and treatment plan. Treatments may include the use of a wart-removal preparation or surgery performed under local anesthesia to safely remove the wart.

### **Wounds and Ulcers**

Foot ulcerations or open wounds are a condition where there is a breakdown in many layers of skin and tissue, sometimes going all the way to the bone. They can be caused by pressure to a weight bearing point on the foot, but they can occur on top of the foot due to pressure from shoes or a bony spur. The risk of ulcer formation is higher in patients with decreased circulation or impaired blood supply to the legs and feet. Diabetics are prone to heel ulcerations, but they can be found in patients with high blood pressure, blood clots, varicose veins, and phlebitis as well.

Treatment for foot ulcerations varies according to the cause of the wound. The goal of the treatment is to close the wound from the inside out as quickly as

## Is Nerve Pain Disrupting Your Lifestyle?

possible. In cases where there is adequate circulation, debridement (removal of dead tissue) may be used around the edges and within the ulcer itself to promote healing. Other treatments include topical treatments with medication, oral medications, compression, taking the pressure off the area (off-loading), and bed rest. New advances have been made in wound care in recent years, including the use of a platelet-derived growth factor as a way to promote healing.



## What Can You Do to Keep Your Feet Healthy?

The first step in dealing with foot pain is to care for your feet. Your feet are the hardest working part of your body. They carry you wherever you need to go, whenever you need to go there, and they do it for years and years. In fact, during your lifetime, you will have traveled on your feet the equivalent of three trips around the entire world. You take, on average, 8,000 steps a day, and will walk or run well over 75,000 miles in your lifetime. Caring for your precious feet is the best step to avoid pain, and is a critical component in eliminating it as well.

### **Here are some steps you can take to keep your feet healthy and functioning:**

First and foremost, do not ignore pain in your feet and ankles. Healthy feet don't have persistent pain or skin that looks unusual. If your pain doesn't subside quickly, please contact our office right away. The sooner we can see you and examine your feet, the quicker we can begin to make the corrections needed to get your feet healthy again. So often we see people who have waited a very long time and suffered needlessly – sometimes for years.

Try to develop a habit of always checking your feet. A great time to do this is right after a shower or during a bath. If you start a habit of carefully drying your feet after bathing (pay special attention to the skin between your toes), you can quickly check your feet to see if you notice any changes. If you see nails that look unusual, you might be seeing a potential fungus developing. If your skin is broken, cracked, or an unusual color, you are noticing abnormalities. Finally, if your foot is changing shape, or you observe new growth, you should make an appointment to see us so we can treat these conditions before they progress and become worse.

If you have diabetes, it is especially important to check your feet very regularly and often. In fact, we recommend that you have someone else help you check your feet, because you may not be able to see or, most importantly,

feel problems. Early detection and treatment may avoid potentially serious complications later.

See a podiatrist if you have a problem with your feet. Treating yourself can often cause problems or exacerbate existing problems. If you are diabetic, you are at greater risk for foot problems so be sure to make an appointment to see us at least once a year. We'd love to see you. We are here to help!



## When To Call a Podiatrist

Podiatrists are the experts on feet and ankles, and should be the first doctor you call when you experience any of the symptoms or pains described in this book. Remember that foot, ankle, or heel pain is never normal regardless of age or activity.

Podiatrists treat foot problems for patients of all ages. You should call a podiatrist when you have pain in your feet or in your heels. If you notice that there is any change at all to your heels or the skin on your feet and ankles, you should see the “expert” on your feet. If you suspect you may have lower extremity nerve pain, you should see your podiatrist right away. The sooner you get treatment for this type of condition, the better the outcome.

If you have diabetes or poor circulation, and you develop any abnormal symptoms, you should see us immediately. You are at much greater risk for serious complications. And if you have diabetes, you should see us at least once a year whether or not you have symptoms or pain. Regular checkups are a great way to help keep your feet healthy.

Most importantly, you should **call us first** when you have pain or any issue that affects your feet.

## **What To Do If You Are In Severe Pain**

Pain is our body's way of telling us that something is wrong. And it is usually true that the more severe the pain, the more serious the problem. If you are experiencing severe foot pain, seek treatment from a podiatrist immediately. Podiatrists specialize in dealing with foot problems and are experts in helping treat your symptoms. You can reach our offices at:

**Harvey R. Danciger, DPM**  
**74-000 Country Club Drive, Suite A-2**  
**Palm Desert, CA 92260**  
**(760) 568-0108**

If you are experiencing an emergency, call 911.

## Final Thoughts

I hope the information in this publication has been helpful. My purpose in sharing it with you is to give you the information you need to take the appropriate actions to care for your feet. I hope it helps you understand lower extremity nerve pain and what can be done to treat it and to keep it from returning. I also wrote it to help you see that podiatric medicine has benefited greatly from the advances medical science is making in the treatment of lower extremity nerve pain and overall foot conditions.

I know that making an appointment to see a doctor isn't always the easiest thing to do, but with knowledge and understanding, you can see that, we, podiatrists can offer you many treatments that can greatly improve the quality of your life. Please make an appointment to come see us to discuss how we can help you with any foot problems you may be experiencing.

Dedicated to Your Health,

*Dr. Harvey Danciger*

## Free Copy of Is Nerve Pain Disrupting Your Lifestyle?



If you have any friends or family members who might benefit from the information provided in this book, we would be happy to provide them with their own copy free of charge. Just tell them to visit our website at [www.CoachellaValleyPodiatrist.com](http://www.CoachellaValleyPodiatrist.com) to request it online OR return this form.

**There is no need to destroy this book. Just photocopy this form, give it to your friend and tell them to mail it or fax it to us.** Fax (760) 568-5110 or mail to 74-000 Country Club Drive, Suite A-2, Palm Desert, CA 92260.

**I want a Free Copy of *Is Nerve Pain Disrupting Your Lifestyle?***

Name: \_\_\_\_\_

Address: \_\_\_\_\_

City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Email: \_\_\_\_\_

Phone number: \_\_\_\_\_

# About the Doctor



As founder of Harvey R. Danciger, DPM, Dr. Danciger leads a talented staff. Dr. Danciger believes that any time you have a problem with your feet or ankles you deserve a friendly doctor and staff who carefully listen to you and care about your health. He is committed to ensuring that patients who put their trust in his podiatric practices receive the finest care available anywhere.

Dr. Danciger completed his undergraduate studies at Syracuse University earning a Bachelor of Science degree. He then took his talents to the Pennsylvania College of Podiatric Medicine, now known as Temple University School of Podiatric Medicine, graduating with a Doctorate of Podiatric Medicine. He completed his Post graduate training in Orange County, California.

Dr. Danciger's devotion to his practice has not gone unnoticed by other professionals in the podiatry field. He has been named a Fellow of the American College of Foot and Ankle Surgeons; Fellow of the Academy of Physicians in Wound Healing; and a Fellow of the American Professional Wound Care Association as well as a Diplomate of the American Board of Podiatric Surgery-Board certified in foot and ankle surgery. He is also a member of the Association of Extremity Nerve Surgeons. He has also served as Treasurer and Chief of Podiatry for the Buena Park Medical Center. All of these positions recognize Dr. Danciger for his excellence in podiatry and mastery of his profession.

Dr. Danciger has practiced in Palm Desert, California for 15 years and in Southern California for over 30 years where he is actively involved with several community service organizations. He served as team podiatrist for Western High School in Anaheim and as volunteer podiatrist for the Jim Cook Day of Hope for Diabetes, March of Dimes Walk-A-Thon and the American Diabetes Association Ride & Stride. He was elected and served eight terms as President of the Coachella Valley Podiatric Medical Society from 2007 thru 2015. Dr. Danciger has lectured on a variety of foot and ankle health related topics. He enjoys treating patients of all ages including children, active adults and the elderly.



**Harvey R. Danciger, DPM**

74-000 Country Club Drive, Suite A-2 • Palm Desert, CA 92260

ph (760) 568-0108 • fx (760) 568-5110

[www.CoachellaValleyPodiatrist.com](http://www.CoachellaValleyPodiatrist.com)